

УДК 517.982

L. HALBEISEN\*

## A SET-THEORETIC APPROACH TO COMPLETE MINIMAL SYSTEMS IN BANACH SPACES OF BOUNDED FUNCTIONS

L. Halbeisen. *A set-theoretic approach to complete minimal systems in Banach spaces of bounded functions*, *Matematychni Studii*, **20** (2003) 162–166.

Using independent families from combinatorial set theory, it is shown that for every infinite cardinal  $\kappa$ ,  $\ell_\infty(\kappa)^*$  contains a subspace which is isomorphic to a Hilbert space of dimension  $2^\kappa$ . This provides a new proof for the first step in the construction of complete minimal systems in Banach spaces of bounded functions.

Л. Халбейзен. *Теоретико-множественный подход к полным минимальным системам в банаховых пространствах ограниченных функций* // *Математичні Студії*. – 2003. – Т.20, №2. – С.162–166.

Используя независимые семейства из комбинаторной множественной теории, мы доказываем, что для каждого бесконечного кардинала  $\kappa$   $\ell_\infty(\kappa)^*$  содержит подпространство, изоморфное гильбертовому пространству размерности  $2^\kappa$ . В результате получено новое доказательство для первого шага конструкции полных минимальных систем в банаховых пространствах ограниченных функций.

**1. Introduction.** Let  $X$  be a Banach space and let  $\{x_\lambda : \lambda \in \Lambda\} \subseteq X$  be an arbitrary set of vectors of  $X$ . Let  $[x_\lambda : \lambda \in \Lambda]$  denote the *closure of the linear span* of  $\{x_\lambda : \lambda \in \Lambda\}$ . A set  $\{x_\lambda : \lambda \in \Lambda\} \subseteq X$  is called a *complete system* if  $[x_\lambda : \lambda \in \Lambda] = X$ , and it is called a *minimal system* if for every  $\lambda' \in \Lambda$ ,  $x_{\lambda'} \notin [x_\lambda : \lambda \in \Lambda \setminus \{\lambda'\}]$ . A *complete minimal system*, abbreviated c.m.s., is a complete system which is also minimal.

Using functionals, we can characterize minimal systems (and consequently c.m.s.) also in the following way: Let  $X$  be a Banach space. A pair of sequences  $\{x_\lambda : \lambda \in \Lambda\} \subseteq X$  and  $\{f_\lambda : \lambda \in \Lambda\} \subseteq X^*$  is called a *biorthogonal system* if  $f_{\lambda'}(x_\lambda) = \delta_{\lambda\lambda}'$ . Now, a sequence  $\{x_\lambda : \lambda \in \Lambda\} \subseteq X$  is minimal if and only if there is a sequence  $\{f_\lambda : \lambda \in \Lambda\} \subseteq X^*$  such that the pair  $(\{x_\lambda : \lambda \in \Lambda\}, \{f_\lambda : \lambda \in \Lambda\})$  is a biorthogonal system. A biorthogonal system which corresponds to a complete minimal system is called a *complete biorthogonal system*.

Even though not every Banach space has a c.m.s. (see e.g., [1] or [2]), it is known that  $\ell_\infty$  has a c.m.s.. The first (not completely correct) proof for the existence of a c.m.s. in  $\ell_\infty$  was given by W. Davis and W. Johnson in [3]. Later, B. Godun gave a correct (and slightly easier) proof in [4]. However, the crucial point in both proofs is the following result due to H. Rosenthal (cf. [5, Proposition 3.4]):

\*The research for this paper was resumed during the *Workshop on Set Theory, Topology, and Banach Space Theory*. The workshop took place in June 2003 at *Queen's University Belfast* and was supported by the author's *Nuffield Foundation Grant NAL/00513/G*.

2000 *Mathematics Subject Classification*: 46B15, 03E05, 46B26.

**Proposition 1.** *The space  $\ell_\infty^*$  contains a subspace isomorphic to a Hilbert space of dimension the continuum.*

Let us briefly sketch why Proposition 1 implies the existence of a c.m.s. in  $\ell_\infty$ : Let  $Y \subseteq \ell_\infty$  be isomorphic to a Hilbert space of dimension the continuum. Since  $Y$  is reflexive,  $Y$  is weakly\* closed (cf. e.g., [5, Proposition 1.2]), and therefore,  $(\perp Y)^\perp = Y$ , where  $\perp Y = \{x \in \ell_\infty : \forall y \in Y (y(x) = 0)\}$  and  $(\perp Y)^\perp := \{x^* \in \ell_\infty^* : \forall x \in \perp Y (x^*(x) = 0)\}$ . Thus,  $(\ell_\infty / \perp Y)^*$  is isomorphic to the Hilbert space  $Y$ , which implies that also  $\ell_\infty / \perp Y$  is isomorphic to  $Y$ . Now, following [4], with the orthonormal basis in  $Y$  we can easily construct a c.m.s. in  $\ell_\infty$ . At this point we like to mention that starting with generalized version of Proposition 1 (cf. [5, p. 203, Remark 2]), a similar construction yields a c.m.s. in  $\ell_\infty(\kappa)$  for any infinite cardinal  $\kappa$ .

Rosenthal’s proof of Proposition 1 involves some deep results from functional analysis. On the other hand, from a set-theoretical point of view a c.m.s. in  $\ell_\infty$  is just a set of bounded real-valued sequences, and therefore, it was natural to seek a more combinatorial or set-theoretical proof of Proposition 1 and the aim of this paper is to provide such a proof.

**Acknowledgement.** I like to thank Anatolij Plichko and Stephanie Halbeisen for pointing me out and explaining me the various steps in the construction of a c.m.s. in  $\ell_\infty$ .

**2. Some set theory. 2.1. Set-theoretic terminology.** Our set-theoretical axioms are the axioms of Zermelo and Fraenkel including the Axiom of Choice. All our set-theoretical notations and definitions are standard and can be found in textbooks like [6].

For a set  $x$ , the *cardinality* of  $x$ , denoted by  $|x|$ , is the least ordinal number  $\alpha$  for which there exists a bijection  $f : \alpha \rightarrow x$ ; such an ordinal number  $\alpha$  is called a *cardinal number* (or just a *cardinal*). The least infinite ordinal number, which is also a cardinal, is denoted by  $\omega$ , thus,  $|\omega| = \omega$ . In particular,  $\omega = \{0, 1, 2, \dots\}$  is the set of natural numbers. A set  $x$  is called finite, if  $|x| \in \omega$ , otherwise it is called infinite. Further it is called countable, if  $|x| \leq \omega$ . For a set  $x$ ,  $\mathcal{P}(x)$  denotes the power set of  $x$  and  $[x]^{<\omega}$  denotes the set of all finite subsets of  $x$ . For a cardinal  $\kappa$ ,  $|\mathcal{P}(\kappa)|$  is denoted by  $2^\kappa$ . For example there exists a bijection between the reals  $\mathbb{R}$  and  $\mathcal{P}(\omega)$ , hence  $|\mathbb{R}| = |\mathcal{P}(\omega)| = 2^\omega$ . For every infinite cardinal we have  $2^\kappa > \kappa$  and  $|\kappa^{<\omega}| = \kappa$ .

**2.2. Independent families.** Let  $\kappa$  be an infinite cardinal and let  $\mathcal{I} \subseteq \mathcal{P}(\kappa)$ , then  $\mathcal{I}$  is called an *independent family* (on  $\kappa$ ), if whenever  $m$  and  $n-1$  belong to  $\omega$ , and  $x_0, \dots, x_m, \dots, x_{m+n}$  are distinct members of  $\mathcal{I}$ , then

$$\left| \bigcap_{0 \leq i \leq m} x_i \setminus \bigcup_{1 \leq j \leq n} x_{m+j} \right| = \kappa.$$

To make this paper self-contained, let us prove the following result due to F. Hausdorff (cf. [7]):

**Proposition 2.** *For any infinite cardinal  $\kappa$ , there is an independent family on  $\kappa$  of cardinality  $2^\kappa$ .*

*Proof.* We just follow Exercise (A6) on p. 288 of [6]. Let

$$J = \{ \langle s, A \rangle : s \subseteq \kappa \text{ and } |s| < \omega \text{ and } A \subseteq \mathcal{P}(s) \}.$$

Notice that  $|J| = \kappa$ , so, it is enough to construct an independent family of cardinality  $2^\kappa$  on  $J$ . For  $x' \subseteq \kappa$ , let  $x := \{ \langle s, A \rangle \in J : x' \cap s \in A \}$ . Then  $\mathcal{I} = \{ x : x' \in \mathcal{P}(\kappa) \}$  is

an independent family on  $J$  of cardinality  $2^\kappa$ . Indeed, let  $x'_0, \dots, x'_m, \dots, x'_{m+n}$  be distinct members of  $\mathcal{P}(\kappa)$  (for some  $m$  and  $n-1$  in  $\omega$ ). Then there is a finite set  $s \subseteq \kappa$  such that for all  $i, j$  with  $0 \leq i < j \leq m+n$  we have  $x'_i \cap s \neq x'_j \cap s$ . Let  $A = \{s \cap x'_i : 0 \leq i \leq m\} \subseteq \mathcal{P}(s)$ , and for every  $\alpha \in \kappa \setminus s$ , let  $s_\alpha = s \cup \{\alpha\}$  and  $A_\alpha = A \cup \{t \cup \{\alpha\} : t \in A\}$ . Then

$$\{\langle s_\alpha, A_\alpha \rangle : \alpha \in \kappa \setminus s\} \subseteq \bigcap_{0 \leq i \leq m} x_i \setminus \bigcup_{1 \leq j \leq n} x_{m+j},$$

which implies that  $\left| \bigcap_{i \leq m} x_i \setminus \bigcup_{m < j \leq n} x_j \right| = \kappa$ , and therefore,  $\mathcal{I}$  is an independent family on  $J$  of cardinality  $2^\kappa$ .  $\square$

As an easy consequence we get the following

**Fact.** If  $\mathcal{I} = \{x_\alpha : \alpha \in 2^\kappa\}$  is an independent family on  $\kappa$  and  $\alpha_1, \dots, \alpha_n$  are finitely many distinct elements of  $2^\kappa$ , then  $\left| \bigcap_{1 \leq i \leq n} y_{\alpha_i} \right| = \kappa$ , where for every  $1 \leq i \leq n$ , the set  $y_{\alpha_i}$  is either equal to the set  $x_{\alpha_i}$  or to its complement  $\kappa \setminus x_{\alpha_i}$ .

**2.3. The Banach spaces  $\ell_2(\kappa)$  and  $\ell_\infty(\kappa)$ .** Let  $\kappa$  be an infinite cardinal. The Banach space  $\ell_\infty(\kappa)$  is the set of all bounded functions from  $\kappa$  to  $\mathbb{R}$ , where for  $x \in \ell_\infty(\kappa)$ ,  $\|x\| = \sup\{x(\alpha) : \alpha \in \kappa\}$ . The Banach space  $\ell_2(\kappa)$  is the set of all functions  $x$  from  $\kappa$  to  $\mathbb{R}$  such that  $\sum_{\alpha \in \kappa} x(\alpha)^2 =: \|x\|^2 < \infty$ . It is common to write  $\ell_2$  and  $\ell_\infty$  instead of  $\ell_2(\omega)$  and  $\ell_\infty(\omega)$  respectively. Like for  $\ell_2$  and  $\ell_\infty$ , one can show that  $\ell_2(\kappa)^* = \ell_2(\kappa)$  and that  $\ell_\infty(\kappa)^*$  is isometric to the space of all finitely additive signed measures  $\mu$  of bounded variation on  $\mathcal{P}(\kappa)$ , supplied with the norm  $\|\mu\| = |\mu|(\kappa)$ , where  $|\mu|$  is the total variation of  $\mu$ .

For  $\alpha, \beta \in \kappa$ , let

$$\delta_\alpha^\beta = \begin{cases} 1 & \text{if } \alpha = \beta, \\ 0 & \text{otherwise,} \end{cases}$$

and let  $e_\alpha : \kappa \rightarrow \{0, 1\}$  be such that  $e_\alpha(\beta) = \delta_\alpha^\beta$ . It is easy to see that the set of vectors  $\{e_\alpha : \alpha \in \kappa\}$  is a c.m.s. of  $\ell_2(\kappa)$ . On the other hand, the set  $\{e_\alpha : \alpha \in \kappa\}$  is much too small to be a c.m.s. of  $\ell_\infty(\kappa)$ . In general, the cardinality of a complete minimal system  $S$  of an infinite dimensional real Banach space  $X$  is always equal to the density character of  $X$ . Indeed, on the one hand, the set of all finite linear combinations of  $S$  with rational coefficients is dense in  $X$ , and on the other hand,  $S$  is discrete in  $X$ . In particular, the density character of  $\ell_\infty(\kappa)$  is  $2^\kappa$ , so, any c.m.s. of  $\ell_\infty(\kappa)$  must have cardinality  $2^\kappa$ .

**3.  $\ell_\infty(\kappa)^*$  contains an isomorphic copy of  $\ell_2(2^\kappa)$ .** Now we are ready to prove the main result.

**Theorem.** *Let  $\kappa$  be an infinite cardinal. Then any independent family on  $\kappa$  of cardinality  $2^\kappa$  induces a subspace of  $\ell_\infty(\kappa)^*$  which is isomorphic to the Hilbert space  $\ell_2(2^\kappa)$ .*

*Proof.* Let  $\mathcal{I} = \{x_\alpha : \alpha \in 2^\kappa\}$  be an independent family on  $\kappa$  of cardinality  $2^\kappa$  (which exists by Proposition 2). Define a measure  $\hat{\mu}$  on the set  $B$  of all Boolean combinations of elements of  $\mathcal{I}$  by stipulating

- $\hat{\mu}(x_\alpha) = \hat{\mu}(\kappa \setminus x_\alpha) = 1/2$  (for all  $x_\alpha \in \mathcal{I}$ ),
- $\hat{\mu}(x_\alpha \cap x_\beta) = \hat{\mu}(x_\alpha \cap (\kappa \setminus x_\beta)) = 1/4$  (for all distinct  $x_\alpha, x_\beta \in \mathcal{I}$ ),

and in general, if  $\alpha_1, \dots, \alpha_n$  are finitely many distinct elements of  $2^\kappa$  and  $0 \leq j \leq n$ , then

$$\hat{\mu} \left( \bigcap_{1 \leq i \leq j} x_{\alpha_i} \cap \bigcap_{j < i \leq n} (\kappa \setminus x_{\alpha_i}) \right) = 2^{-n}.$$

The measure  $\hat{\mu}$  induces a normalized linear functional  $\varphi_{\hat{\mu}}$  on a subspace of  $\ell_\infty(\kappa)$ . Thus, by the normed space version of the Hahn-Banach Extension Theorem, there is a normalized functional on all of  $\ell_\infty(\kappa)$  which extends the functional  $\varphi_{\hat{\mu}}$ . In particular, there is a measure  $\mu$  on  $\mathcal{P}(\kappa)$  with  $\|\mu\| = 1$ , such that  $\mu|_B \equiv \hat{\mu}$ . For every  $\alpha \in 2^\kappa$  let  $f_\alpha: \kappa \rightarrow \{1, -1\}$  such that

$$f_\alpha(\lambda) = \begin{cases} 1 & \text{if } \lambda \in x_\alpha, \\ -1 & \text{otherwise.} \end{cases}$$

Now, for every  $\alpha \in 2^\kappa$ , let the measure  $\mu_\alpha$  on  $\mathcal{P}(\kappa)$  be defined by

$$\mu_\alpha(E) = \mu(E \cap x_\alpha) - \mu(E \cap (\kappa \setminus x_\alpha)),$$

and let  $\varphi_\alpha$  be the linear functional on  $\ell_\infty(\kappa)$  induced by the measure  $\mu_\alpha$ . It is not hard to see that for all  $\alpha, \beta \in 2^\kappa$ ,  $\varphi_\alpha(f_\beta) = \delta_\alpha^\beta$  and that  $\|\varphi_\alpha\|_{\ell_\infty(\kappa)^*} = 1$ . Let  $Y = [\varphi_\alpha : \alpha \in 2^\kappa] \subseteq \ell_\infty(\kappa)^*$ , and let  $\sum_{i=1}^n a_i \varphi_{\alpha_i} \in Y$ .

**Claim.** For each  $\varepsilon = (\varepsilon_1, \dots, \varepsilon_n) \in \{-1, 1\}^n$  we have

$$\left\| \sum_{i=1}^n a_i \varphi_{\alpha_i} \right\|_{\ell_\infty(\kappa)^*} = 2^{-n} \sum_{\varepsilon \in \{-1, 1\}^n} \left| \sum_{i=1}^n \varepsilon_i a_i \right|.$$

*Proof.* For each  $\varepsilon = (\varepsilon_1, \dots, \varepsilon_n) \in \{-1, 1\}^n$  let  $E_\varepsilon = \bigcap_{1 \leq i \leq n} y_{\alpha_i}$ , where

$$y_{\alpha_i} = \begin{cases} x_{\alpha_i} & \text{if } \varepsilon_i a_i \geq 0, \\ \kappa \setminus x_{\alpha_i} & \text{otherwise.} \end{cases}$$

By the fact mentioned above,  $|E_\varepsilon| = \kappa$ , and by the properties of the measure  $\mu$  we get  $\mu(E) = 2^{-n}$ . Notice that for any distinct  $\varepsilon$  and  $\varepsilon'$  in  $\{-1, 1\}^n$  we have  $E_\varepsilon \cap E_{\varepsilon'} = \emptyset$  and that  $\kappa = \bigcup_{\varepsilon \in \{-1, 1\}^n} E_\varepsilon$ . Further, for every  $\varepsilon \in \{-1, 1\}^n$  let  $f_\varepsilon: \kappa \rightarrow \{\pm 1, 0\}$  be such that

$$f_\varepsilon(\lambda) = \begin{cases} 1 & \text{if } \lambda \in E_\varepsilon \text{ and } |\varepsilon_1 a_1 + \dots + \varepsilon_n a_n| \geq 0, \\ -1 & \text{if } \lambda \in E_\varepsilon \text{ and } |\varepsilon_1 a_1 + \dots + \varepsilon_n a_n| < 0, \\ 0 & \text{otherwise,} \end{cases}$$

and let  $f = \sum_{\varepsilon \in \{-1, 1\}^n} f_\varepsilon$ . It is not hard to verify that for each  $\varepsilon \in \{-1, 1\}^n$  we have

$$(a_1 \varphi_{\alpha_1} + \dots + a_n \varphi_{\alpha_n})(f_\varepsilon) = 2^{-n} |\varepsilon_1 a_1 + \dots + \varepsilon_n a_n|,$$

and therefore,

$$\left( \sum_{i=1}^n a_i \varphi_{\alpha_i} \right)(f) = 2^{-n} \sum_{\varepsilon \in \{-1, 1\}^n} \left| \sum_{i=1}^n \varepsilon_i a_i \right|.$$

Now, since the  $E_\varepsilon$ 's are pairwise disjoint,  $\|f\|_{\ell_\infty(\kappa)} = 1$ , and by the construction of  $f$  we finally get

$$\left\| \sum_{i=1}^n a_i \varphi_{\alpha_i} \right\|_{\ell_\infty(\kappa)^*} = 2^{-n} \sum_{\varepsilon \in \{-1, 1\}^n} \left| \sum_{i=1}^n \varepsilon_i a_i \right|.$$

□

Hence, by Khintchine's inequality, there is a constant  $c = 1/\sqrt{2}$  such that

$$c \cdot \sqrt{\sum_{i=1}^n a_i^2} \leq \left\| \sum_{i=1}^n a_i \varphi_{\alpha_i} \right\|_{\ell_\infty(\kappa)^*} \leq \sqrt{\sum_{i=1}^n a_i^2},$$

which implies that the space  $Y \subseteq \ell_\infty(\kappa)^*$  is isomorphic to the Hilbert space  $\ell_2(2^\kappa)$  and completes the proof.  $\square$

**Remark.** For an infinite cardinal  $\kappa$ , the Banach space  $c_0(\kappa)$  is the set of all functions  $x$  from  $\kappa$  to  $\mathbb{R}$  such that for every  $\varepsilon > 0$ , the set  $\{\alpha < \kappa : |x(\alpha)| > \varepsilon\}$  is finite. Now, the Theorem admits the following generalization: *Let  $\kappa$  be an infinite cardinal. Then the space  $(\ell_\infty(\kappa)/c_0(\kappa))^*$  contains a subspace which is isomorphic to  $\ell_2(2^\kappa)$ . Consequently we get: For every infinite cardinal  $\kappa$ , the space  $\ell_\infty(\kappa)/c_0(\kappa)$  has a c.m.s..*

## REFERENCES

1. Plichko A. *A Banach space without a fundamental biorthogonal system*, Doklady Akademii Nauk SSSR, **254** (1980), №4 798–801 (in Russian); translation in Soviet Mathematical Doklady, **22** (1980), №2, 450–453.
2. Godun B. V., Kadets M. I. *Banach spaces without complete minimal systems*, Funktsional'nyi Analiz i Ego Prilozheniya **14** (1980), №4, 67–68 (in Russian); translation in Functional Analysis and its Applications, **14** (1980), №4, 301–302.
3. Davis W. J., Johnson W. B. *On the existence of fundamental and total bounded biorthogonal systems in Banach spaces* Studia Mathematica, **45** (1973), 173–179.
4. Godun B. V. *Complete biorthogonal systems in Banach spaces*, Funktsional'nyi Analiz i Ego Prilozheniya, **17** (1983), №1, 1–7 (in Russian); translation in Functional Analysis and its Applications **17** (1983), №1, 1–5.
5. Rosenthal H. P. *On quasi-complemented subspaces of Banach spaces, with an appendix on compactness of operators from  $L^p(\mu)$  to  $L^r(\mu)$* , Journal of Functional Analysis, **4** (1969), 176–214.
6. Kunen K. *Set Theory, an Introduction to Independence Proofs*, North Holland, Amsterdam (1983).
7. Hausdorff F. *Über zwei Sätze von G. Fichtenholz und L. Kantorovitch*, Studia Mathematica, **6** (1936), 18–19.

Department of Pure Mathematics, Queen's University Belfast,  
Belfast BT7 1NN, Northern Ireland  
halbeis@qub.ac.uk

*Received 19.09.2003*